

WHAT IS CLAIMED IS:

1. A composition useful for determining the presence of vitamin B12 in a sample, the composition comprising an antibody that is capable of specifically binding to intrinsic factor only in the absence of vitamin B12, and that is released from binding in the presence, and upon the binding, of vitamin B12 to intrinsic factor.

2. A kit for determining the presence of vitamin B12 in a sample, the kit comprising intrinsic factor and labelled antibody, the antibody being one which will specifically bind to the intrinsic factor in an amount related to the amount of vitamin B12 present in the sample, wherein the antibody is capable of binding to intrinsic factor only in the absence of vitamin B12, and is released from binding in the presence, and upon the binding, of vitamin B12 to intrinsic factor.

3. The kit of claim 2 further comprising a second antibody which will specifically bind to intrinsic factor without regard to the presence or absence of vitamin B12.

4. The kit of claim 3 wherein the second antibody is bound to a solid phase support.

5. A kit for assaying for vitamin B12 in a sample comprising (a) solid phase support to which is bound a predetermined amount of an antibody that is capable of specifically binding to intrinsic factor only in the absence of vitamin B12, and that is released from binding in the presence, and upon the binding, of vitamin B12 to intrinsic factor, and (b) a predetermined amount of a labelled intrinsic factor.

6. The kit of claim 5 wherein the label is alkaline phosphatase and further comprising a substrate to detect the presence or amount of the label.

7. A method of obtaining a monoclonal antibody capable of binding to intrinsic factor only in the absence of vitamin B12, and being released from binding in the presence, and upon the binding, of vitamin B12 to intrinsic factor, comprising the steps of:

- a) immunizing an animal with substantially purified intrinsic factor;
- b) isolating splenic lymphocytes from the immunized animal;
- c) fusing the isolated splenic lymphocytes with a plasmocytoma cell line to obtain a plurality of hybridoma clones which secrete antibody;
- d) extracting free vitamin B12 from a predetermined amount of culture supernatant containing antibody from each hybridoma clone;
- e) contacting a first sample of each extracted antibody-containing supernatant with intrinsic factor in the presence of vitamin B12;
- f) contacting a second sample of each extracted antibody-containing supernatant with intrinsic factor in the absence of vitamin B12;
- g) contacting an enzyme labelled antibody which specifically binds to immunoglobulin with each of the first and second samples;

- h) detecting the presence of labelled antibody present in each of the first and second samples; and
- i) isolating the hybridomas which secrete antibodies which bound to intrinsic factor only in the absence of vitamin B12.

5
8g. The method of claim 8 wherein the extraction step (d) is performed using dextran coated charcoal.

Rule
1.126
910. A diagnostic assay method for vitamin B12 in a liquid sample comprising:

- 10 (a) contacting the sample with a known amount of labelled intrinsic factor and a known amount of an antibody bound to a solid phase which specifically binds to intrinsic factor in an amount related to the presence or amount of vitamin B12 in the sample, said antibody being capable of binding to a site on intrinsic factor that is distinct from the
- 15 site on intrinsic factor to which vitamin B12 binds and which binds to intrinsic factor only in the absence of vitamin B12, and that is released from of B12? binding in the presence, wherein the intrinsic
- 20 factor will specifically bind to vitamin B12 in the sample to form a vitamin B12-intrinsic factor complex;

- (b) separating the vitamin B12-intrinsic factor complex; and
- (c) determining the amount of vitamin B12 by measuring the amount of label associated with the vitamin B12-intrinsic factor complex or the amount label bound to the antibody on the solid phase.

5
10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995

10 11. The method of claim 10 wherein the antibody is a monoclonal antibody.